American Journal of Artificial Intelligence and Intelligent Systems
ISSN: - applied

Integrated Remote Sensing and Machine Learning Frameworks for Shallow Water Bathymetry Retrieval: From Empirical Models to Attention-Based Intelligence Systems

[1] Dr. Alejandro Martín-Ruiz
[1] Department of Geomatics and Environmental Engineering, University of Barcelona, Spain

Received: 10th Nov 2025 | Received Revised Version: 18th Nov 2025 | Accepted: 25th Nov 2025 | Published: 02th Dec 2025

Volume 01 Issue 02 2025 | Crossref DOI: 10.64917/ajaiis/V01I02-01

Abstract

Accurate estimation of shallow water bathymetry is a long-standing scientific and practical challenge with profound implications for coastal management, marine navigation, ecological monitoring, disaster mitigation, and sustainable development. Traditional bathymetric survey methods, while highly precise, are constrained by cost, spatial coverage, and operational complexity, particularly in shallow, turbid, or ecologically sensitive waters. Consequently, satellite-derived bathymetry has emerged as a critical alternative, leveraging multispectral remote sensing data to infer water depth based on light–water interactions. Over the past several decades, research in this domain has evolved from simple empirical and semi-empirical models toward sophisticated machine learning and deep learning approaches capable of capturing complex nonlinear relationships between spectral signals and underwater topography.

This study presents a comprehensive, theoretically grounded research article synthesizing empirical remote sensing bathymetry techniques with contemporary machine learning and attention-based intelligence models. Drawing strictly on the provided literature, the article develops an integrated analytical framework that connects early correlation-based fathoming models, multispectral depth inversion techniques, and comparative multi-source remote sensing studies with modern supervised learning algorithms such as random forests, support vector regression, artificial neural networks, convolutional neural networks, and transformer-based attention mechanisms. By doing so, the research situates bathymetry inversion within a broader computational intelligence paradigm, highlighting how methodological advances in machine learning—originally developed in fields such as legal judgment prediction and natural language processing—offer conceptual and practical insights applicable to marine remote sensing.

The methodology section elaborates in depth on the theoretical foundations of satellite-derived bathymetry, including radiative transfer principles, spectral attenuation behavior, and the assumptions underlying empirical and semi-analytical inversion models. It then extends this discussion to machine learning methodologies, explaining how feature representation, nonlinear regression, ensemble learning, kernel-based methods, and hierarchical neural architectures improve depth retrieval accuracy and robustness. Particular emphasis is placed on attention mechanisms and residual learning, not as direct bathymetric tools, but as transferable conceptual models that inspire adaptive weighting of spectral information and context-aware depth estimation.

The results are presented through descriptive analytical synthesis rather than numerical tabulation, demonstrating consistent trends across studies: machine learning-assisted bathymetry generally outperforms traditional empirical approaches in heterogeneous coastal environments, while deep learning models show superior generalization when sufficient training data are available. The discussion critically evaluates limitations related to data dependency, model interpretability, transferability across regions, and environmental variability, while also outlining future research directions that integrate explainable artificial intelligence, multi-source data fusion, and attention-based architectures tailored specifically for marine environments.

Overall, this article contributes a unified theoretical narrative that bridges marine remote sensing and modern artificial intelligence research. By articulating the conceptual continuity between empirical bathymetric inversion and attention-driven learning systems, it provides a foundation for future interdisciplinary research aimed at developing accurate, scalable, and interpretable satellite-derived bathymetry solutions.

Keywords: Satellite-derived bathymetry, multispectral remote sensing, machine learning, convolutional neural networks, attention mechanisms, shallow water mapping.

© **2025** Dr. Alejandro Martín-Ruiz. This work is licensed under a Creative Commons Attribution 4.0 International License (CC BY 4.0). The authors retain copyright and allow others to share, adapt, or redistribute the work with proper attribution.

Cite This Article: Martín-Ruiz, D. A. (2025). Integrated Remote Sensing and Machine Learning Frameworks for Shallow Water Bathymetry Retrieval: From Empirical Models to Attention-Based Intelligence Systems. American Journal of Artificial Intelligence and Intelligent Systems, 1(2), 1-5. https://doi.org/10.64917/ajaiis/V01I02-01

1. Introduction

The measurement and representation of underwater topography, commonly referred to as bathymetry, constitute a foundational element of marine science and coastal engineering. Bathymetric information underpins a wide range of applications, including nautical charting, harbor and port management, sediment transport analysis, habitat mapping, coastal erosion assessment, and disaster risk reduction related to tsunamis and storm surges. Historically, bathymetric data acquisition has relied on in situ survey techniques such as lead-line sounding, single-beam and multi-beam echo sounding, and airborne LiDAR bathymetry. While these methods provide high accuracy, they are constrained by financial cost, logistical complexity, temporal inflexibility, and limited spatial coverage, particularly in shallow or environmentally sensitive coastal waters.

The advent of satellite remote sensing introduced a transformative paradigm for bathymetric estimation, enabling large-scale, repetitive, and cost-effective observation of marine environments. Early research demonstrated that multispectral satellite imagery contains exploitable information related to water depth, particularly in optically shallow waters where sunlight penetrates the water column and reflects from the seabed. The attenuation of light at different wavelengths, modulated by water depth, turbidity, and bottom type, forms the physical basis of satellite-derived bathymetry. Empirical and semi-empirical models were initially developed to exploit these relationships, correlating spectral reflectance values with known depth measurements to generate predictive models (Zhang et al., 1998; Dang and Ding, 2003).

Over time, researchers recognized that simplistic linear or log-linear relationships were insufficient to capture the complexity of coastal and shallow marine environments. Variability in water quality, bottom albedo, and atmospheric conditions introduced nonlinearities that challenged traditional regression-based approaches. This recognition led to the exploration of more flexible modeling techniques, including artificial neural networks, support vector regression, and ensemble learning methods (Wang and Zhang, 2005; Wang et al., 2018; Qiu et al., 2019). These machine learning approaches offered the ability to model complex, nonlinear interactions without explicit physical parameterization, thereby improving depth inversion accuracy in heterogeneous conditions.

Concurrently, advancements in computer science and artificial intelligence—particularly in machine learning, deep learning, and attention-based architectures—have reshaped multiple research domains. Notably, attention mechanisms and transformer models have revolutionized natural language processing and legal analytics by enabling models to dynamically weight contextual information and capture long-range dependencies (Vaswani et al., 2023; Sukanya and Priyadarshini, 2021). While these techniques were not originally designed for remote sensing or bathymetry, their underlying principles of adaptive feature weighting, hierarchical representation learning, and contextual awareness hold significant conceptual relevance for multispectral depth estimation.

Despite the rapid methodological evolution, the literature reveals a fragmentation between traditional remote sensing bathymetry research and broader machine learning advancements. Many studies focus narrowly on algorithmic performance within specific case studies, without situating their contributions within a unified theoretical framework that spans empirical modeling, machine learning regression, and attention-based intelligence. Furthermore, comparative analyses across different methodological paradigms often lack depth in theoretical justification, focusing instead on numerical accuracy metrics without fully exploring the epistemological implications of model choice.

This article addresses these gaps by developing a comprehensive, theory-driven research narrative that

integrates empirical satellite-derived bathymetry models with contemporary machine learning and attention-based approaches. By drawing exclusively on the provided references, the study ensures conceptual coherence and scholarly rigor while offering extensive elaboration on the theoretical underpinnings, methodological evolution, and interpretive challenges of bathymetric inversion. The objective is not merely to summarize existing work, but to critically synthesize it into an integrated framework that highlights continuity, divergence, and future potential at the intersection of marine remote sensing and artificial intelligence.

2. Methodology

The methodological foundation of this research is rooted in analytical synthesis rather than experimental replication. Given the diverse nature of the referenced studies, which span empirical remote sensing models, machine learning regression techniques, and attention-based intelligence systems, the methodology adopted here involves a structured, theory-oriented examination of underlying principles, algorithmic mechanisms, and inferential assumptions. This approach enables a deep exploration of methodological evolution while maintaining fidelity to the original studies.

At the core of satellite-derived bathymetry lies the interaction between electromagnetic radiation and the aquatic environment. Multispectral sensors capture reflected sunlight across visible and near-infrared bands, with shorter wavelengths such as blue and green exhibiting greater penetration into clear water. Early empirical models, such as those developed for the South China Sea, established statistical relationships between spectral reflectance ratios and measured water depths (Dang and Ding, 2003). These models often relied on linear or logarithmic transformations to stabilize variance and enhance correlation, implicitly assuming homogeneous water properties and bottom reflectance.

Subsequent research expanded these approaches by incorporating multi-source remote sensing data and comparative inversion techniques. By analyzing data from different sensors and spectral configurations, researchers demonstrated that depth inversion accuracy is sensitive to spatial resolution, spectral bandwidth, and sensor-specific noise characteristics (Lu et al., 2019). Methodologically, this stage marked a shift from single-model reliance toward comparative evaluation, emphasizing robustness and transferability across datasets.

The introduction of machine learning represented a methodological inflection point. Artificial neural networks, particularly backpropagation-based models, were among the earliest nonlinear techniques applied to bathymetry retrieval. These models treated spectral reflectance values as input features and water depth as the output target, learning complex mappings through iterative weight optimization (Wang and Zhang, 2005). Unlike empirical regression, neural networks did not require explicit functional forms, allowing them to adapt to nonlinearities inherent in coastal environments.

Support vector regression further enriched the methodological toolkit by introducing kernel-based learning and margin optimization. By mapping input features into high-dimensional spaces, support vector regression models captured subtle nonlinear patterns while maintaining strong generalization properties (Wang et al., 2018). The incorporation of fuzzy membership concepts addressed uncertainty in training data, reflecting the recognition that bathymetric measurements and spectral observations are subject to noise and ambiguity.

Ensemble learning methods, particularly random forest algorithms, represented another methodological advancement. By aggregating predictions from multiple decision trees, random forests reduced variance and improved robustness against overfitting (Qiu et al., 2019). The theoretical appeal of ensemble methods lies in their ability to approximate complex functions through the collective behavior of simpler models, an idea that resonates with earlier empirical approaches while offering greater flexibility.

Deep learning architectures, especially convolutional neural networks, further transformed bathymetric inversion methodology. By leveraging spatial context and hierarchical feature extraction, convolutional models moved beyond pixel-wise regression toward spatially aware depth estimation (Ai et al., 2020). Methodologically, this shift acknowledged that water depth is not solely a function of local spectral values, but also of spatial patterns related to seabed morphology and coastal processes.

While attention mechanisms and transformer models were developed primarily for sequential data in natural language processing, their methodological principles offer transferable insights. Attention-based models dynamically weight input features based on contextual relevance, enabling selective focus on informative components (Vaswani et al., 2023). In the context of bathymetry, this concept translates into adaptive weighting of spectral bands,

spatial neighborhoods, or temporal observations, even if explicit transformer implementations are not yet widespread in marine remote sensing.

By synthesizing these methodological strands, this study adopts a holistic analytical framework that views bathymetric inversion as an evolving continuum of representational complexity. Each methodological layer builds upon its predecessors, retaining core physical intuition while embracing increasing computational sophistication.

3. Results

The results of this research are articulated through descriptive synthesis rather than quantitative comparison, reflecting the integrative and theoretical nature of the study. Across the referenced literature, several consistent patterns emerge regarding the performance and implications of different bathymetric inversion methodologies.

Empirical and semi-empirical models demonstrate reliable performance in optically clear, shallow waters where assumptions of uniform water properties hold. Studies conducted in regions such as the South China Sea and the southern Huanghai Sea illustrate that carefully calibrated empirical relationships can yield acceptable depth estimates, particularly for operational applications requiring rapid assessment (Dang and Ding, 2003; Zhang et al., 2009). However, these models exhibit sensitivity to environmental variability and often require region-specific recalibration.

Machine learning models consistently outperform traditional empirical approaches in heterogeneous environments. Neural network-based models show enhanced capacity to capture nonlinear spectral-depth relationships, leading to improved accuracy across varying bottom types and water conditions (Wang and Zhang, 2005). Support vector regression and random forest models further enhance robustness by balancing flexibility with generalization, reducing susceptibility to overfitting and noise (Wang et al., 2018; Qiu et al., 2019).

Deep learning approaches, particularly convolutional neural networks, achieve the highest reported performance levels when sufficient training data are available. Their ability to integrate spatial context results in smoother and more realistic bathymetric surfaces, aligning closely with known geomorphological features (Ai et al., 2020). However, these gains come at the cost of increased computational complexity and reduced interpretability.

The conceptual integration of attention mechanisms, while not directly evaluated in bathymetric studies, yields important inferential results. Research in legal judgment prediction and computational intelligence demonstrates that attention-based models excel at identifying salient features and contextual dependencies in complex datasets (Sukanya and Priyadarshini, 2021; Chen et al., 2022). By analogy, these results suggest that attention-inspired strategies could enhance bathymetric inversion by dynamically prioritizing informative spectral and spatial cues.

4. Discussion

The synthesized results underscore a fundamental trade-off in satellite-derived bathymetry between physical interpretability and predictive flexibility. Empirical models offer transparency and ease of implementation, grounded in physical intuition about light attenuation. However, their rigidity limits applicability in complex or changing environments. Machine learning and deep learning models address this limitation by embracing data-driven representation, but they introduce challenges related to interpretability, data dependency, and transferability.

One critical limitation across advanced models is their reliance on high-quality training data. Bathymetric ground truth is often sparse or unevenly distributed, constraining model generalization. Moreover, deep learning models function as black boxes, complicating scientific interpretation and stakeholder trust. Attention mechanisms partially mitigate this issue by offering insight into feature importance, suggesting a pathway toward more explainable bathymetric models.

Future research should prioritize the integration of physical knowledge with data-driven learning, developing hybrid models that embed radiative transfer constraints within machine learning architectures. Additionally, multi-source data fusion, incorporating optical, radar, and LiDAR observations, holds promise for enhancing robustness. Attention-based architectures tailored to multispectral and spatial data represent a particularly compelling avenue, drawing on successful applications in legal analytics and computational intelligence.

5. Conclusion

This article has presented an extensive, theory-driven synthesis of satellite-derived bathymetry research, tracing its evolution from empirical remote sensing models to advanced machine learning and attention-based intelligence systems. By integrating insights from marine science and artificial intelligence, the study demonstrates that bathymetric inversion is not merely a technical problem, but

a paradigmatic example of interdisciplinary convergence. The future of shallow water bathymetry lies in models that balance physical understanding, computational power, and interpretability, ensuring accurate and sustainable mapping of the world's coastal environments.

References

1. Ai, B., Wen, Z., Wang, Z., Wang, R., Su, D., Li, C., & Yang, F. (2020). Convolutional neural network to retrieve water depth in marine shallow water area from remote sensing images. IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing, 13, 2888–2898.
2. Aletras, N., Tsarapatsanis, D., Preoţiuc-Pietro, D., & Lampos, V. (2016). Predicting judicial decisions of the European Court of Human Rights: A natural language processing perspective. PeerJ Computer Science, 2, e93.
3. Ao, Y., Li, H., Zhu, L., Ali, S., & Yang, Z. (2018). The linear random forest algorithm and its advantages in machine learning assisted logging regression modeling. Journal of Petroleum Science and Engineering, 174, 776–789.
4. Casal, G., Monteys, X., Hedley, J., Harris, P., Cahalane, C., & McCarthy, T. (2019). Assessment of empirical algorithms for bathymetry extraction using Sentinel-2 data. International Journal of Remote Sensing, 40, 2855–2879.
5. Chen, J., Du, L., Liu, M., & Zhou, X. (2022). Mulan: A multiple residual article-wise attention network for legal judgment prediction. ACM Transactions on Asian and Low-Resource Language Information Processing, 21, 1–15.
6. Cheng, K., & Zhu, Y. (2007). A summary of machine learning and related algorithms. Statistics and Information Forum, 22, 105–112.
7. Dang, F., & Ding, Q. (2003). A technique for extracting water depth information from multispectral scanner data in the South China Sea. Marine Science Bulletin, 22, 55–60.
8. Lu, T., Chen, S., Tu, Y., Yu, Y., Cao, Y., & Jiang, D. (2019). Comparative study on coastal depth inversion based on multi-source remote sensing data. Chinese Geographical Science, 29, 192–201.
9. Qiu, Y., Shen, W., Hui, X., & Zhang, H. (2019). Satellite-derived bathymetry using random forest model. Journal of Ocean Technology, 34, 75–79.
10. Shang, X. (2022). A computational intelligence model for legal prediction and decision support. Computational Intelligence and Neuroscience, 2022, 1–8.
11. Sukanya, G., & Priyadarshini, J. (2021). A meta analysis of attention models on legal judgment prediction system. International Journal of Advanced Computer Science and Applications, 12.
12. Sukanya, G., & Priyadarshini, J. (2024). Analysis on word embedding and classifier models in legal analytics. AIP Conference Proceedings, 2802, 140001.
13. Vaswani, A., Jones, L., Shazeer, N., Parmar, N., Uszkoreit, J., Gomez, A. N., Kaiser, Ł., & Polosukhin, I. (2023). Attention is all you need. arXiv preprint arXiv:1706.03762.
14. Wang, J., Ma, Y., & Zhang, J. (2018). Multiple kernel support vector regression based on fuzzy membership for remote sensing water depth fusion detection. Marine Environmental Science, 37, 130–136.
15. Wang, Y., & Zhang, Y. (2005). Study on remote sensing of water depth based on BP artificial neural networks. Ocean Engineering, 23, 37–42.
16. Zhang, D., Zhang, Y., & Wang, W. (1998). Establishment of a RS-fathoming correlation model. Journal of Hohai University, 26, 98–102.
17. Zhang, Y., Zhang, Y., Zhang, D., & Qian, Y. (2009). An underwater bathymetry reversion in the radial sand ridge group region of the southern Huanghai Sea using the remote sensing technology. Acta Oceanologica Sinica, 31, 39–45.